

Hemelron

Premium source for iron deficiency



- Bioavailable polypeptide form
- Easily absorbed







newrootsherbal.com

Hemelron



Iron deficiency causes a form of anemia, leading to clinically low levels of hemoglobin (red blood cells). Iron deficiency anemia can cause fatigue, shortness of breath, dizziness, headaches, and lack of stamina. Persons at risk include women of reproductive age during menstruation, people with celiac and inflammatory bowel disease, and conditions that contribute to nutrient malabsorption.

Highly bioavailable **Heme**lron is isolated from animal sources for maximum intestinal absorption. It's the particular type of iron that is easily recognized and exploited by the body to form the structural scaffolding for hemoglobin. Hemoglobin is the specialized protein within blood that binds to oxygen within the lungs for use in cellular respiration throughout the entire body. **Heme**Iron also does not lead to common side effects associated with elemental forms of iron that include: stomach upset, diarrhea, and constipation.

New Roots Herbal's **Heme**lron is formulated with a full spectrum of complementary nutrients. It contains vitamin C to enhance intestinal iron absorption, with the biologically active forms of folic acid (methyl folate) and vitamin B₁₂ (methylcobalamin) included as critical cofactors for hemoglobin synthesis.

Each vegetable capsule contains:

Iron (heme iron polypeptide from porcine hemoglobin)	11 mg
Iron (heme iron polypeptide from porcine hemoglobin) Vitamin C Folate (from calcium L-5-methyltetrahydrofolate) 3: Vitamin B ₁₂ (methylcobalamin). 3:	90 mg
Folate (from calcium L-5-methyltetrahydrofolate)	33 mcg
Vitamin B ₁₂ (methylcobalamin)	33 mcg

Other ingredients: Vegetable magnesium stearate, microcrystalline cellulose, and silicon dioxide in a non-GMO vegetable capsule composed of vegetable carbohydrate gum and purified water. NPN 80061737 · V0514-R2

Adults: Take 1 capsule one to three times daily with a full glass of water, with or without food, or as directed by your health-care practitioner. Do not lie down for 30 minutes after taking this product. If you are taking other medications, take this product a few hours before or after them. Consult a health-care practitioner for use beyond 7 weeks. Consult a health-care practitioner to monitor blood iron content.

Manufactured under strict GMP (Good Manufacturing Practices).

